

# $\eta$ -Simple semigroups without zero and $\eta^*$ -simple semigroups with a least non-zero idempotent

Roman S. Gigoń

Received: 16 September 2011 / Accepted: 12 May 2012 / Published online: 31 May 2012  
© The Author(s) 2012. This article is published with open access at Springerlink.com

**Abstract** A semigroup  $S$  is called  $\eta$ -simple if  $S$  has no semilattice congruences except  $S \times S$ . Tamura in (Semigroup Forum 24:77–82, 1982) studied  $\eta$ -simple semigroups with a unique idempotent. In the present paper we consider a more general situation, that is, we investigate  $\eta$ -simple semigroups (without zero) with a least idempotent. Moreover, we study  $\eta^*$ -simple semigroups with zero which contain a least non-zero idempotent.

**Keywords**  $\eta$ -Simple semigroup · Least idempotent ·  $E$ -Inversive semigroup

## 1 Preliminaries

Let  $S$  be a semigroup and  $a \in S$ . An element  $x \in S$  is called a *weak inverse* of  $a$  if  $xax = x$ ; the set of all weak inverses of  $a$  is denoted by  $W_S(a)$ . A semigroup  $S$  is said to be  *$E$ -inversive* if for every  $a \in S$  there is  $x \in S$  such that  $ax \in E_S$ , where  $E_S$  (or briefly  $E$ ) is the set of idempotents of  $S$  (more generally, if  $A \subseteq S$ , then  $E_A$  denotes the set of idempotents of  $A$ ). If  $A \subseteq S$ , then by  $A^*$  we shall mean the set of all non-zero elements of  $A$ . Since each semigroup with zero is  $E$ -inversive, then we define a semigroup  $S$  with zero to be  *$E$ -inversive* if for all  $a \in S^*$  there exists  $x \in S$  such that  $ax \in E_S^*$ . Finally, put  $W_S^*(a) = W_S(a) \setminus \{0\}$  ( $a \in S$ ). Recall from [3] that a semigroup  $S$  [with zero] is  $E^{[*]}$ -inversive if and only if  $W_S^{[*]}(a) \neq \emptyset$  for every  $a \in S^{[*]}$ .

**Lemma 1.1** *A semigroup  $S$  [with zero] is  $E^{[*]}$ -inversive if and only if every [non-zero] ideal of  $S$  contains some [non-zero] idempotent of  $S$ .*

---

Communicated by Marcel Jackson.

R.S. Gigoń (✉)

Institute of Mathematics and Computer Science, Wrocław University of Technology,  
Wyb. Wyspiańskiego 27, 50-370 Wrocław, Poland  
e-mail: [romekgigon@tlen.pl](mailto:romekgigon@tlen.pl)

*Proof* Suppose that every non-zero ideal of  $S$  contains some non-zero idempotent of  $S$ ,  $a \in S^*$ . Then  $S^1 a S^1$  contains at least one non-zero idempotent of  $S$ , that is,  $xay = e$  for some  $x, y \in S^1$  (in fact, we may suppose that  $x, y \in S$ ),  $e \in E_S^*$ . Hence  $exaye = e$ , so  $(yex)a(yex) = yex \neq 0$ ; otherwise  $0 = xa(yex) = (xay)ex = ex$ . Thus  $0 = exay = e$ , a contradiction. Consequently,  $yex \in W_S^*(a)$ .

The converse implication is clear.  $\square$

**Lemma 1.2** *Let  $S$  be an  $E^{[*]}$ -inversive semigroup. Then  $eSe$  is  $E^{[*]}$ -inversive for every  $e \in E_S^{[*]}$ .*

*Proof* Observe first that  $e \in eSe$ , so  $eSe \neq \{0\}$ . Let  $a \in (eSe)^*$  and  $x \in W_S^*(a)$ . Then  $x = xax = x(eae)x$ . Hence  $exe = (exe)a(exe)$ . Furthermore, if  $exe = 0$ , then we get  $xe = [(xe)a(ex)]e = (xea)(exe) = 0$ , so  $x = (xe)a(ex) = 0$ , a contradiction. Thus  $exe \in W_{eSe}^*(a)$ , as exactly required.  $\square$

We say that a semigroup  $S$  is a *semilattice* if  $a^2 = a$ ,  $ab = ba$  for all  $a, b \in S$ . Further, a congruence  $\rho$  on a semigroup  $S$  is called a *semilattice congruence* if  $S/\rho$  is a semilattice. It is clear that the least semilattice congruence  $\eta$  on an arbitrary semigroup exists. Finally, a semigroup is said to be  $\eta$ -simple if  $\eta = S \times S$ .

The next lemma follows immediately from the Second Isomorphism Theorem.

**Lemma 1.3** *A homomorphic image of an  $\eta$ -simple semigroup is  $\eta$ -simple.*

Let  $S$  be a semigroup. Recall that the *natural partial order* is the relation  $\leq$ , defined on  $E_S$  by  $e \leq f$  if  $e = ef = fe$ . We say that a semigroup  $S$  (without zero) has a least idempotent  $e$  if  $e \leq f$  for every  $f \in E_S$ . Note that If  $S$  has a zero, say  $0$ , then clearly  $0$  is the least element of  $E_S$  with respect to  $\leq$ , but in such a case, we may say that  $S$  has a *least non-zero idempotent* if  $E_S^*$  contains the least element with respect to the natural partial order.

Let  $A$  be an ideal of a semigroup  $S$ . We say that  $S$  is an *ideal extension* of the semigroup  $A$  by the semigroup  $T$  if the Rees semigroup  $S/A$  is isomorphic to  $T$ . Finally, an ideal  $P$  of a semigroup  $S$  is called *prime* if the condition  $ab \in P$  implies that  $a \in P$  or  $b \in P$  for all  $a, b \in S$ .

## 2 The main results

Remark that by Corollary 3.9 of [4], a semigroup  $S$  is  $\eta$ -simple if and only if  $S$  has no proper prime ideals.

**Proposition 2.1** *Let  $S \neq S^0$  be an  $\eta$ -simple semigroup with a least idempotent. Then  $S$  is  $E$ -inversive. Moreover,  $S$  is an ideal extension of a group by an  $\eta$ -simple semigroup.*

*Proof* Let  $e$  be the least element of  $E_S$ . Then every ideal of  $S$  must contain  $e$ . Indeed, suppose by way of contradiction that there is an ideal  $A$  of  $S$  such that  $e \notin A$ . Let  $B$

be the set theoretic union of all such ideals  $A$  of  $S$ . Then clearly  $B$  is the largest ideal of  $S$  such that  $e \notin B$ . Next, consider the Rees quotient  $S/B$ . Notice that we may think about  $S/B$  as a semigroup with zero, where all products not falling in  $S/B$  are zero. Consider now an arbitrary non-zero ideal  $C$  of  $S/B$ . Then by construction of  $B$ ,  $\{e\}$  must belong to  $C$ . Hence the intersection of all non-zero ideals of  $S/B$  contains  $\{e\}$ . In particular,  $S/B$  is  $E^*$ -inversive (see Lemma 1.1). Also,  $B$  is a prime ideal of  $S$ . Indeed, let  $a, b \notin B$  be such that  $ab \in B$ . Then  $fg \in B$  for some  $f, g \in E_S \setminus B$  (because  $S/B$  is  $E^*$ -inversive). Hence  $e = efg \in B$  (which is a contradiction). It follows that  $S$  has a proper semilattice congruence (by the above remark), a contradiction with the assumption of the theorem. Consequently, every ideal of  $S$  must contain  $e$ . Thus  $S$  has a kernel  $G$  (say) and  $S$  is  $E$ -inversive (Lemma 1.1). Hence for every  $a \in S$  there exists  $x \in S$  such that  $ax, xa \in E_S$ . Therefore  $e = (ax)e = a(xe) \in aS$ . We may equally well show that  $e \in Sa$ . It follows easily that  $S$  contains both a minimum left ideal  $L$  (say) and a minimum right ideal  $R$  (say). Furthermore, for every  $a \in S$ ,  $La$  is a minimal left ideal of  $S$  (see [1], Lemma 2.32). Hence  $La = L$ , so  $L$  is an ideal of  $S$  (and  $L = L^2$ ). We can show in a similar way that  $R$  is an ideal of  $S$ , so  $L = R = G = eS = Se$  (because  $Se \subseteq L, eS \subseteq R$ , since  $e \in L, R$ ). Consequently,  $G = eSe$ . Indeed, evidently  $eSe \subseteq SeS = G$ . Also,  $G = GG = eSSe \subseteq eSe$ . By Lemma 1.2,  $G$  is an  $E$ -inversive monoid (with an identity element  $e$ ). Moreover, if  $f \in E_{eSe}$ , then  $fe = ef = f$  i.e.  $f \leq e$ . Thus  $f = e$ . Consequently,  $G$  is a group ideal of  $S$  and so  $S$  is an ideal extension of the group  $G$  by the semigroup  $S/G$  which is  $\eta$ -simple, by Lemma 1.3, as required.  $\square$

**Lemma 2.2** *Let  $S \neq S^0$  be an  $\eta$ -simple semigroup with the least idempotent  $e$ . Then  $ea = ae$  for all  $a \in S$ .*

*Proof* Let  $a \in S$ . Then  $ea, ae \in eSe = eS = Se$ , where  $eSe$  is a group ideal of  $S$  (see the proof of Proposition 2.1). Hence  $e \cdot ae = ae, ea \cdot e = ea$ . Thus  $ea = ae$ .  $\square$

A congruence on a semigroup is called a *group congruence* if the quotient semigroup is a group.

**Corollary 2.3** *Let  $S \neq S^0$  be an  $\eta$ -simple semigroup with a least idempotent, say  $e$ . Then the mapping  $s \rightarrow es$  of  $S$  onto the group  $eS$  is an epimorphism leaving the elements of  $eS$  fixed. Moreover, the congruence  $\sigma$  induced by this morphism, that is  $\sigma = \{(a, b) \in S \times S : ea = eb\}$ , is the least group congruence on  $S$ .*

*Proof* The first part of the result follows from Proposition 2.1 and Lemma 2.2. Further, if  $\rho$  is a group congruence on  $S$ , then clearly  $(s, es) \in \rho$  for every  $s \in S$ . Hence  $\sigma \subseteq \rho$ .  $\square$

**Remark 1** Notice that if a semigroup  $S \neq S^0$  with the least idempotent  $e$  is  $\eta$ -simple, then  $\rho_{eS} \cap \sigma = 1_S$  and so  $S$  is a subdirect product of an ( $E$ -inversive)  $\eta$ -simple semigroup  $S/eS$  (with zero) and the group  $eS$ .

Further, the converse of Proposition 2.1 is valid.

**Theorem 2.4** *A semigroup  $S$  without zero is  $\eta$ -simple and has a least idempotent if and only if it is an ( $E$ -inversive) ideal extension of a group by an  $\eta$ -simple semi-group.*

*Proof* The direct part follows from Proposition 2.1.

Conversely, let  $G$  be a group ideal of  $S$  (with an identity  $e$ ) and  $a \in S$ . Then  $ea \in G$ , say  $ea = g$ . It follows that  $g^{-1}ea = e \in Sa$ . We may equally well show that  $e \in aS$ , so  $e$  is the least idempotent of  $E_S$ . Further, if  $\rho$  is a semilattice congruence on  $S$ , then (according to the proof of Theorem 5 in [5])  $\rho \cap (G \times G) = G \times G$ . It follows that  $\rho_G \subset \rho$ , where  $\rho_G$  is the Rees congruence on  $S$  modulo  $G$ . Hence there is an epimorphism of  $S/\rho_G$  onto  $S/\rho$ . In fact, this morphism induced on  $S/\rho_G$  a semi-lattice congruence. Since  $S/\rho_G$  is  $\eta$ -simple, then  $S/\rho$  must be trivial. Consequently,  $\rho = S \times S$ , as exactly required.  $\square$

Remark that if a semigroup  $S$  is a left [right] group (i.e.  $S \times S = \mathcal{L}[\mathcal{R}]$ ), then  $S$  is  $\eta$ -simple. Indeed, let  $S$  be a left [right] group. Then  $S \times S = \mathcal{L}[\mathcal{R}] \subseteq \mathcal{J} \subseteq \eta$ .

**Theorem 2.5** *A semigroup  $S$  without zero is  $\eta$ -simple and has an idempotent  $e$  such that  $ef = e$  [ $fe = e$ ] if and only if it is an ( $E$ -inversive) ideal extension of a left [right] group by an  $\eta$ -simple semigroup.*

*Proof* ( $\implies$ ). Let  $ef = e$  for every  $f \in E_S$ . We may equally well show like above (see the proof of Proposition 2.1) that  $e$  belongs to every ideal of  $S$ . Hence  $S$  has a kernel, say  $K$ . In particular,  $S$  is  $E$ -inversive. It follows that  $e \in Sa$  for every  $a \in S$ . Thus  $S$  contains a minimum left ideal  $L$  and  $L = La$  for all  $a \in S$  (so  $L = L^2$ ). Therefore  $K = Se$  is a left simple semigroup (by Theorem 2.35 [1]) and so  $K$  is a left group (by the dual of Theorem 1.27 [1]). Consequently,  $S$  is an ideal extension of the left group  $K$  by the semigroup  $S/K$  which is  $\eta$ -simple.

( $\impliedby$ ). Let  $K$  be a left group ideal of  $S$ ,  $e \in E_K$  and  $a \in S$ . Then  $ea \in K$ , say  $ea = k$ . It follows that  $ek^{-1}ea = ek^{-1}k = e \in Sa$ , where  $k^{-1}$  is some inverse of  $k$  in  $K$  (since  $E_K$  is a left zero semigroup). Hence if  $f \in E_S$ , then  $e = sf$  for some  $s \in S$ . Thus  $ef = e$ . We have just shown that  $ef = e$  for all  $e \in E_K$ ,  $f \in E_S$ . Further, if  $\rho$  is a semilattice congruence on  $S$ , then  $\rho \cap (K \times K) = K \times K$  (by the preceding remark) and so  $\rho = S \times S$  (by the proof of Theorem 2.4).  $\square$

**Corollary 2.6** *Let  $S$  be a simple semigroup. If  $S$  has an idempotent  $e$  such that  $ef = e$  [ $fe = e$ ] for every  $f \in E_S$ , then  $S$  is a left [right] group.*

*Proof* Indeed, in such a case,  $\mathcal{J} = S \times S$ . It is almost evident (and also well-known) that  $\mathcal{J} \subseteq \eta$ . Hence  $S$  is  $\eta$ -simple, so  $S$  contains a left [right] group ideal  $K$ . Thus  $S = K$ .  $\square$

Notice that if  $S$  is a completely simple semigroup (see [2], Sect. 3.2), then the Green's relation  $\mathcal{H}$  is a band congruence on  $S$  (see Lemma III.2.4 in [2]). Further, every left [right] group  $S$  is completely simple and  $E_S$  is a left [right] zero semigroup. It follows, from the above, that if  $S$  is a left [right] group, then  $S/\mathcal{H}$  is a left [right] zero semigroup.

A semigroup  $S$  is said to be *congruence-free* if it has exactly two congruences.

**Proposition 2.7** *Let  $S$  be a congruence-free semigroup without zero. If  $S$  has an idempotent  $e$  such that  $ef = e$  [ $fe = e$ ] for every  $f \in E_S$ , then  $S$  is a simple group.*

*Proof* Let  $ef = e$  for every  $f \in E_S$ . Since  $S$  is congruence-free, then either  $\eta$  is the identity or the universal relation on  $S$ . In the former case,  $S$  is a semilattice, but then  $e$  is the zero of  $S$ , a contradiction with the assumption of the proposition. It follows that  $S$  is  $\eta$ -simple. By Theorem 2.5,  $S$  contains a left group ideal  $K$ . Hence  $S$  is itself a left group. From the above remark we conclude that either  $\mathcal{H} = 1_S$  or  $\mathcal{H} = S \times S$ . In the former case,  $S$  must be a left zero semigroup. Since  $|S| > 1$ , then the partition  $\{\{e\}, S \setminus \{e\}\}$  of  $S$  induced a proper congruence on  $S$ , a contradiction. Thus  $\mathcal{H} = S \times S$ , so  $E_S = \{e\}$ , since  $\mathcal{H}$  separates idempotents of  $S$ . Consequently,  $S$  is a simple group.  $\square$

Next, consider a semigroup  $S$  with zero such that the set  $E_S^*$  contains a least idempotent, say  $e$ . Remark that  $fg \neq 0$  for all  $f, g \in E_S^*$  (in fact, if  $e \in E_S^*$  has the property that  $ef = e$  [ $fe = e$ ] for every  $f \in E_S^*$ , then also  $gh \neq 0$  for all  $g, h \in E_S^*$ ).

Since a semigroup with zero adjoined has a proper semilattice congruence, then we shall say that a semigroup with zero is  $\eta^*$ -simple if  $S$  has at most two semilattice congruences, namely: (i)  $S \times S$  or (ii) the congruence induced by the partition  $\{\{0\}, S^*\}$ . Clearly, the partition  $\{\{0\}, S^*\}$  of a semigroup  $S$  with zero induces a semilattice congruence on  $S$  if and only if  $S$  is a semigroup with zero adjoined.

Recall that a semigroup  $S$  with zero is called a 0-group if  $S^*$  is a group.

**Theorem 2.8** *A semigroup  $S$  with zero is  $\eta^*$ -simple and has a least non-zero idempotent if and only if it is an  $E^*$ -inversive semigroup with zero adjoined (and so  $S^*$  is an  $E$ -inversive semigroup with a least idempotent) and it is an ideal extension of a 0-group by an  $\eta$ -simple semigroup.*

*Proof* ( $\Rightarrow$ ). Let  $e$  be a least non-zero idempotent of  $S$ . We can show that every non-zero ideal of  $S$  contains  $e$  (see the proof of Proposition 2.1 and the above remark). In particular,  $S$  is  $E^*$ -inversive (Lemma 1.1). Hence for every  $a \in S^*$  there is  $x$  such that  $xa$  is a non-zero idempotent of  $S$ . Thus  $e \in Sa$ . We may equally well show that  $e \in aS$ . Next, if  $a, b \in S^*$ , then (by the above)  $e = xa, e = by$  for some  $x, y \in S$ . Hence  $e = x(ab)y$  and so  $ab \in S^*$ . Consequently,  $S$  has no proper zero divisors. Thus  $S^*$  is an  $E$ -inversive semigroup with a least idempotent  $e$  and so  $S^*$  is an ideal extension of a group  $G$  by an  $\eta$ -simple semigroup (Theorem 2.4). It follows that  $S$  is an ideal extension of a 0-group  $G^0$  by an  $\eta$ -simple semigroup. Indeed,  $S/G^0$  must have a proper zero divisor (otherwise  $G^0$  is a non-zero prime ideal of  $S$ ).

The opposite implication follows easily from the proof of Theorem 2.4.  $\square$

A non-zero [left [right]] ideal  $A$  of a semigroup  $S$  with zero is called 0-minimum if it is contained in every non-zero [left [right]] ideal of  $S$ .

Further, a semigroup  $S$  with zero is called *categorical* if  $abc = 0$  implies that either  $ab = 0$  or  $bc = 0$  for all  $a, b, c \in S$ .

Finally, we have the following theorem.

**Theorem 2.9** *Let  $S$  be a categorical semigroup (with zero). Then  $S$  is  $\eta^*$ -simple and has a non-zero idempotent  $e$  such that  $ef = e$  [ $fe = e$ ] for every  $f \in E_S^*$  if and only if it is an  $E^*$ -inversive semigroup with zero adjoined (and so  $S^*$  is an  $E$ -inversive semigroup with a least idempotent) and it is an ideal extension of a left [right] group with zero adjoined by an  $\eta$ -simple semigroup.*

*Proof* ( $\implies$ ). Let  $ef = e$  for every  $f \in E_S^*$ . We may equally well show like above that  $e$  belongs to every non-zero ideal of  $S$ . Hence  $S$  has a 0-minimum ideal  $K$ . In particular,  $S$  is  $E^*$ -inversive. It follows that  $e \in Sa$  for all  $a \in S^*$ . Thus  $S$  contains a 0-minimum left ideal  $L$  (and  $L = Le$ , so  $L = L^2$ ). Therefore  $K = Se$  is a left 0-simple semigroup (by Theorem 2.35 in [1]), so  $K^*$  is a left simple semigroup (Theorem 2.27 in [1]). Thus  $K^*$  is a left group (by the dual of Theorem 1.27 in [1]). Further, suppose that  $ea = 0$  for some  $a \in S$  and let  $b \in S^*$ . Then  $e = sb$  for some  $s \in S$ . Hence  $sba = 0$ . Thus  $ba = 0$  (since  $S$  is categorical), so  $\{0, a\}$  is a left ideal of  $S$ . It follows that either  $\{0, a\} = K$  or  $a = 0$ . Consequently,  $ea \neq 0$  for all  $a \in S^*$ . Therefore  $ab \neq 0$  for all  $a, b \in S^*$ . Indeed, if  $ab = 0$  for some  $a, b \in S^*$ , then  $eb = 0$ , a contradiction from the above. We conclude that  $S^*$  is an  $E$ -inversive semigroup, so  $S^*$  is an ideal extension of the left group  $K^*$  by the semigroup  $S/K$  which is  $\eta$ -simple (Theorem 2.5). Hence  $S$  is an ideal extension of the left group  $K$  with zero adjoined by the semigroup  $S/K$  which is  $\eta$ -simple, since  $K$  is not a prime ideal of  $S$ .

The opposite implication follows from the proof of Theorem 2.5.  $\square$

**Open Access** This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

## References

1. Clifford, A.H., Preston, G.B.: The Algebraic Theory of Semigroups, Vol. I. Math. Surveys, vol. 7. Am. Math. Soc., Providence (1961)
2. Howie, J.M.: An Introduction to Semigroup Theory. Academic Press, London (1984)
3. Mitsch, H., Petrich, M.: Basic properties on  $E$ -inversive semigroups. Commun. Algebra **28**, 5169–5182 (2000)
4. Petrich, M.: The maximal semilattice decomposition of a semigroup. Math. Z. **55**, 68–82 (1964)
5. Tamura, T.: Semilattice indecomposable semigroups with a unique idempotent. Semigroup Forum **24**, 77–82 (1982)